

## SP 2/0 Host Cell Proteins

### Western Blot Kit for the Detection of SP 2/0 Host Cell Proteins Catalog # F185

#### Intended Use

This kit is intended for use in determining the presence of Host Cell Protein contamination in products manufactured by recombinant expression in SP 2/0 host cells. The kit is for **Research and Manufacturing Use Only** and is not intended for diagnostic use in humans or animals.

#### Summary and Explanation

Recombinant expression by the hybridoma SP 2/0 cell line is a relatively simple and cost effective method for production of monoclonal antibodies. Many of these recombinantly produced antibodies are intended for use as therapeutic agents in humans and animals and as such must be highly purified. The manufacturing and purification process of these products leaves the potential for contamination by host cell proteins from SP 2/0. Such contamination can result in adverse toxic or immunological reactions and thus it is desirable to reduce host cell contamination to the lowest levels practical.

The Western blot technique is a common analytical tool used to detect host cell protein contamination. Samples to be evaluated are first subjected to polyacrylamide gel electrophoresis (PAGE) often in the presence of detergent such as SDS and a reducing agent such as dithiothreitol (DTT). Under these conditions, proteins will migrate through the gel and be separated as a function of their mass and charge. In the Western Blot procedure the proteins which are separated on the gel are then electrophoretically transferred to a membrane, typically made of nitrocellulose or polyvinylidene difluoride (PVDF) where these proteins are essentially irreversibly adsorbed onto the membrane. After a blocking step with an irrelevant protein such as bovine serum albumin to saturate unoccupied adsorption sites on the membrane, the membrane is then exposed to a solution containing goat antibodies to SP 2/0 which have been labeled with the enzyme Horse Radish Peroxidase (HRP). These antibodies will in turn bind to any transferred proteins for which they are specific. After a wash step to remove any non-immunologically bound antibody, the membrane is finally exposed to the substrate, Tetramethylbenzidine (TMB) specifically formulated to precipitate on the membrane. Those locations where the enzyme labeled antibody has bound to a transferred protein will be indicated by the generation of a substrate chromogen product in characteristic bands on the membrane. In this way, specific components in a complex mixture of proteins can be conclusively identified.

The antibodies used in this kit are polyclonal and were generated by a proprietary procedure designed to elicit a very broad reactivity to a large number of SP 2/0 antigens. These antibodies have been shown to react to more than 40 different SP 2/0 protein bands from SDS/DTT solubilized SP 2/0 cells, or

from HCP's found in conditioned SP 2/0 culture media after one dimensional PAGE separation. This kit provides a simple, very sensitive system capable of detecting as little as 1ng of protein per band. As such this kit can be used as a process development tool or routine quality control method to monitor the optimal removal of host cell contaminants. For more sensitive detection of SP 2/0 HCP's in downstream or final product it is recommended to use an ELISA. In addition, to almost 2 log orders of improved sensitivity over typical Western Blots, an ELISA is more quantitative. *Cygnus Technologies* also has available an SP 2/0 HCP ELISA kit, Cat. No. F180.

#### Reagents & Materials Provided

Component	Product #
<b>Anti-SP 2/0:HRP Conjugate</b> Affinity purified goat antibody conjugated to HRP in a protein matrix with preservative. 2x50mL	F187
<b>SP 2/0 Control Antigen</b> Solubilized and diluted SP 2/0 cell proteins with preservative. 1x50µL	F186
<b>TMB Substrate</b> 3,3',5,5' Tetramethylbenzidine. 1x100mL	F129
<b>Block/Wash Concentrate (20X)</b> Tris buffered saline with bovine serum albumin and preservative. 1x50 mL	F062

#### Storage & Stability

\* All reagents should be stored at 2°C to 8°C for stability until the expiration date printed. **DO NOT FREEZE.**

\* Reconstituted wash solution is stable until the expiration date of the kit.

#### Materials Required But Not Provided

Blotting/Transfer membranes (nitrocellulose or PVDF)  
Distilled water  
1-liter container for wash solution storage  
Reagent trays

#### Precautions

\* For Research or Manufacturing use only.

\* At the concentrations used in this kit, none of the reagents are believed to be harmful.

\* This kit should only be used by qualified technicians.

## Preparation of Reagents

\* Prior to PAGE, the Control Antigen (#F186), should be diluted in the same running buffer (reducing or non-reducing) as the samples.

\* Bring all reagents to room temperature.

\* Dilute wash concentrate to 1 liter in distilled water, label with kit lot and expiration date, and store at 4°C.

## Procedural Note

Complete washing of the membrane to remove excess unreacted goat SP 2/0:HRP is essential to minimize background color and achieve maximum sensitivity.

## Limitations

1. The antibodies were generated against a SP 2/0 cells commonly used in recombinant procedures. A typical SDS/DTT solubilized preparation of SP 2/0 can show more than 40 distinct bands. However, there can be no guarantee that this assay will detect all proteins or protein fragments from SP 2/0.

2. Typical Western Blot sensitivity limits for detection of SP 2/0 proteins are approximately 1ng per band. The detection limits for some bands could be higher than 1ng per band.

3. It is recommended that other methods of host cell protein contamination be evaluated to ensure the absence of significant contamination. *Cygnus Technologies* also manufactures a microtiter plate based ELISA for SP 2/0, Cat. No. F180. This kit is approximately 100 fold more sensitive than the Western Blot and as such is more suitable for final product testing where HCP levels will typically be below the sensitivity limits of Western Blot.

## Blotting Protocol Guidelines

\* Optimization of the conditions for the PAGE and electrophoretic transfer to the membrane needs to be experimentally determined by each user in order to achieve maximum sensitivity for the Western Blotting procedure.

\* The following procedure is typical of one which might be used to give satisfactory results on 8x10cm mini-gels. This procedure is offered as an example only. You may find it advantageous to vary reagent volumes, antibody:enzyme conjugate dilution, incubation times and washing steps to achieve the desired results.

\* The SP 2/0 Control Antigen is a diluted extract which is provided to serve as a positive control for the entire procedure from electrophoresis to completion of the blotting protocol. This material should be treated in the same way as samples, i.e. dilution in reducing or non-reducing PAGE running buffers. Recommended final dilution of the control antigen is 1:1 to 1:5. The development of at least 20 bands during the substrate step is indicative of a satisfactory run.

## Typical Protocol for Minigel (8x10cm) Blots

1. After electrophoretic transfer from the PAGE gels onto the membrane, place the membrane into 40mL of diluted Block/Wash solution in an appropriately sized reagent tray. Allow the blocking of the membrane to proceed for 30 minutes with agitation or rotation to ensure good mixing and even diffusion through the membrane.

2. Pour off the Block/Wash solution and add 20mL\* of Anti-SP 2/0:HRP Conjugate (#F187). Incubate with gentle agitation for 2 hours at room temperature.

3. Carefully pick up the membrane by the corner using forceps. Touch off any drops of the antibody:enzyme conjugate and transfer to a clean reagent tray containing 40mL of Block/Wash solution. Allow the membrane to wash for 5 minutes with agitation. Pour off the Block/Wash solution and replace with another 40mL. Repeat for a total of 4 washes.

4. Transfer the membrane to a clean reagent tray containing 20mL of the TMB substrate (#F129). Incubate with gentle agitation for approximately 30 minutes.\*\*

5. Stop the substrate by rinsing the membrane in distilled water.

\* The reagent tray should be a length and width such that the volume of enzyme conjugate added will completely cover the membrane and allow for free flowing of the solution around the membrane.

\*\* The point at which to stop the substrate incubation should be determined by the user for each blot. The reaction should be stopped before the background color becomes so intense that there is insufficient contrast between positive bands and background. In some cases sensitivity can be increased by incubating with the anti-SP 2/0:HRP for up to 16 hours. If it is necessary to stop the substrate reaction much earlier than 30 minutes the user may consider diluting the antibody:enzyme conjugate in Block/Wash Solution or shortening the immunological incubation step to less than 2 hours.

## Ordering Information/ Customer Service

To place an order or to obtain additional product information contact *Cygnus Technologies*:

[www.cygnustechnologies.com](http://www.cygnustechnologies.com)

Cygnus Technologies, Inc.  
4701 Southport Supply Rd. SE, Suite 7  
Southport, NC 28461 USA  
Tel: 910-454-9442  
Fax: 910-454-9443  
Email: [techsupport@cygnustechnologies.com](mailto:techsupport@cygnustechnologies.com)